



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ :

C07D 451/02, A61K 51/02, 101/02, 31/41

A1

(11) International Publication Number:

WO 95/11901

(43) International Publication Date:

4 May 1995 (04.05.95)

(21) International Application Number: PCT/US94/12225

(22) International Filing Date: 24 October 1994 (24.10.94)

(30) Priority Data:

08/142,584

25 October 1993 (25.10.93)

US

(71) Applicants: THE GENERAL HOSPITAL CORPORATION [US/US]; 55 Fruit Street, Boston, MA 02114 (US). THE PRESIDENT AND FELLOWS OF HARVARD COLLEGE [US/US]; 17 Quincy Street, Cambridge, MA 02138 (US). ORGANIX, INC. [US/US]; 65 Cummings Park, Woburn, MA 01801 (US). NORTHEASTERN UNIVERSITY [US/US]; 360 Huntington Avenue, Boston, MA 02115 (US).

(72) Inventors: ELMALEH, David, R.; 38 Hartman Road, Newton, MA 02159 (US). MADRAS, Bertha, K.; 32 Montrose Street, Newton, MA 02158 (US). MELTZER, Peter; 8 Rumford Road, Lexington, MA 02173 (US). HANSON, Robert, N.; 8 Shornecliffe Road, Newton, MA 02158 (US).

(74) Agent: FREEMAN, John, W.; Fish & Richardson, 225 Franklin Street, Boston, MA 02110-2804 (US).


(81) Designated States: European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).

Published

With international search report.

Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.

(54) Title: SUBSTITUTED 2-CARBOXYALKYL-3-(FLUOROPHENYL)-8-(3-HALOPROPEN-2-YL) NORTROPANES AND THEIR USE AS IMAGING AGENTS FOR NEURODEGENERATIVE DISORDERS



(57) Abstract

Disclosed are certain cocaine analogs useful for imaging of cocaine receptors and dopamine receptors. Also disclosed are analogs useful for imaging diagnostics for Parkinson's disease.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

- 1 -

SUBSTITUTED 2-CARBOXYALKYL-3-(FLUOROPHENYL)
-8-(3-HALOPROPEN-2-YL) NORTROPANES AND
THEIR USE AS IMAGING AGENTS
FOR NEURODEGENERATIVE DISORDERS

5

Background of the Invention

The invention relates to substituted nortropanes and their use to image: a) dopamine-transporter-containing neurons, particularly in connection with diagnosis and study of certain neurodegenerative disorders; and b) cocaine receptors.

There is a need for diagnostic agents and markers of neurogenerative disorders such as Parkinson's disease. For example, exclusion at an early stage of Parkinson's disease as the cause of symptoms may be useful information in diagnosing other conditions. Moreover, early diagnosis of Parkinson's disease can facilitate the introduction of putative prophylactic drug therapy (e.g., deprenyl) prior to the onset of more severe symptoms. See, Kaufman and Madras (1991) *Synapse* 9:43-49. Detection of nerve cell depletion in the presymptomatic phase in an animal model of Parkinson's disease would also be useful, e.g., when using the model to evaluate therapies for Parkinson's disease. See, Hantraye et al. (1992) *Neuro Reports* 3:265-268; and Hantraye et al. (1992) *Soc. Neurosci. Abstra.* 18:935.

There is a particular need for diagnostic agents and markers of neurogenerative disorders that selectively target a dopamine transporting protein (the dopamine transporter) in preference to another protein known as the serotonin transporter. In normal brain tissue, the dopamine:serotonin transporter density ratio is approximately 10:1. In certain neurodegenerative disorders, such as Parkinson's disease, nerve cells that produce dopamine (and on which the dopamine transporter is located) undergo severe depletion, while serotonin

- 2 -

nerve cells are less affected. The dopamine:serotonin transporter ratio can fall to 50% in Parkinson's disease.

Various substances (particularly cocaine and cocaine congeners) are potent inhibitors of dopamine transport in the striatum of the brain because they bind to the dopamine transporter. The more strongly these substances block dopamine transport, the more strongly they bind to sites on the dopamine transporter which have been labeled by [³H]cocaine or by a compound known as [³H]CFT (also known as [³H]WIN 35,428).¹ See, Madras et al., (1989) *J. Pharmacol. Exp. Ther.* 251:131-141; and Madras et al. (1989) *Mol. Pharmacol.* 36:518-524.

CFT (WIN 35,428) and similar substances exhibit markedly decreased binding in the Parkinson's diseased brain. See, Madras et al. *Soc. Neurosci. Abst.* 16:14, 1990; and Kaufman and Madras (1991) *Synapse* 9:43-49. The hope that these compounds might be Parkinson's markers is further supported by the parallel between loss of binding and loss of dopamine in the diseased brain (Madras et al. *Catechol. Symp.* 193, 1992).

Dopamine transporter-binding compounds that have been studied include N-allyl-2 carboxymethyl-3 β -fluorophenyltropane. See, Madras et al. (1990) *Pharmacology, Biochemistry & Behavior* 35:949-953; and Milius et al. (1991) *J. Med. Chem.* 34:1728-1731.

Goodman et al. report (1992) *J. Nuclear Med.* 33:890) the synthesis of 2- β -carbomethoxy-3- β -(4-chlorophenyl)-8-(¹²³I-iodopropene-2yl) nortropane. The selectivity of this analog for the dopamine over the serotonin transporter is not discussed.

CFT or WIN 35,428 is approximately 15 times more selective for the dopamine over the serotonin transporter. The 4-chloro analog is 4-fold selective.

¹ 2-carboxymethyl-3-(4-fluorophenyl)tropane.

- 3 -

See Carroll et al., *J. Med. Chem.* 35:969 (1992); and
Meltzer et al., *J. Med. Chem.* 36:855-862 (1993).

Summary of the Invention


We have discovered that fluorination of the phenyl
5 substituent of N-haloallyl phenyl nortropanes yields
compounds that are unusually effective for several
purposes. The 3E and 3Z isomers of 2- β -carbomethoxy-3- β -
(4-fluorophenyl)-8-(3-I-iodopropen-2-yl) nortropene are
particularly preferred. We have designated this
10 preferred compound Iodcaltropene or IACFT. Most
preferably, the compounds according to the invention are
labeled with radionuclides of the halogen series, such as
 ^{123}I or ^{125}I .

The special properties of the claimed compounds
15 are best understood with the following background.
Cocaine recognition sites are localized on the dopamine
transporter, which itself is localized on dopamine nerve
terminals. Drugs that bind to these sites therefore have
potential uses which include: (i) imaging probes for
20 neurodegenerative disorders; and (ii) imaging probes for
dopamine transporter/cocaine binding sites.

Because of the unique anatomical location of the
cocaine recognition sites, a high affinity probe for
imaging of these sites in vivo in the brain can be
25 carried out using PET or SPECT imaging. Such imaging is
useful for diagnosing or monitoring Parkinson's disease,
a neurological disorder characterized by the degeneration
of dopamine nerve terminals or by aging.

- 4 -

Accordingly, one aspect of the invention features a compound of formula:



wherein the following condition is imposed on that formula:

- 5 R is $-\text{CH}_3$, $-\text{CH}_2\text{CH}_3$ (α configuration; β configuration or both), $-\text{CH}(\text{CH}_3)_2$, $-(\text{CH}_2)_n\text{CH}_3$, $-(\text{CH}_2)_nC_6H_4X$, $-C_6H_4X$, $-C_6H_5$, $-OCH_3$, $-OCH_2CH_3$, $-OCH(\text{CH}_3)_2$, $-OC_6H_5$, $-OC_6H_4X$, $-O(\text{CH}_2)_nC_6H_4X$, or $-O(\text{CH}_2)_n\text{CH}_3$; wherein X is $-\text{Br}$, $-\text{Cl}$, $-\text{I}$, $-\text{F}$, $-\text{OH}$, $-OCH_3$, $-\text{CF}_3$, $-\text{NO}_2$, $-\text{NH}_2$, $-\text{CN}$, $-\text{NHCOCH}_3$, $-\text{N}(\text{CH}_3)_2$, $-(\text{CH}_2)_n\text{CH}_3$, CHOCH_3 , or $-\text{C}(\text{CH}_3)_3$ and n is between 0 and 6 inclusive
- 10

A second aspect of the invention features the same formula, except that the substituent on the N-allyl group can generally be any halogen, preferably $-\text{I}$ or $-\text{F}$.

- 15 In a third aspect, the invention features a method of detecting parkinsonism in a human patient, involving administering to the patient a detectably labelled compound of the invention and detecting its binding to CNS tissue, e.g., by quantifying dopamine terminals with

- 5 -

that compound (e.g., using positron emission tomography (PET) or single-photon emission computed tomography (SPECT)).

In a fourth aspect, the compound is used to
5 monitor cocaine binding sites of the CNS (e.g. to determine site occupancy by potential cocaine therapeutics).

Such compounds also may be useful in treatment of neurodegenerative disorders or cocaine abuse.

10 Preferred compounds according to the invention are characterized as follows: a) the 2 substituent is in the β position; b) the 3 substituent is in the β position; c) R is $-O-CH_3$; the 8 substituent is either the E isomer or the Z isomer. More preferably the halo substituent on
15 the N-allyl moiety is $-I$ or $-Br$ (particularly a radionuclide of $-I$ or $-Br$); ^{18}F provides also a useful label. Most preferably, the compound is Iodoaltropine: 2- β -carbomethoxy-3- β -(4-fluorophenyl)-8-(3E-iodopropen-2-yl) nortropine.

20 Also in preferred embodiments, the compound contains a radioactive label (particularly a gamma or position emitter such as ^{123}I , ^{125}I , ^{18}F , or ^{11}C ; ^{123}I is particularly preferred) or a ^{18}F fluoro label as part of the 3-halopropen-2-yl substituent.

Therapeutic compositions according to the invention comprise a compound as described above formulated in a pharmaceutically acceptable carrier. Such compositions can be used to selectively image cocaine binding regions of the central nervous system of a human patient by administering detectably labelled compound to the central nervous system and detecting the binding of that compound to CNS tissue. The compositions can also be used to detect parkinsonism in a human patient by administering detectably labelled compound to the patient and detecting the binding of the compound to

- 6 -

CNS tissue (e.g., by position emission tomography (PET) or by single-photon emission computed tomography (SPECT)). Such a compound also may be useful in treatment of neurodegenerative disorders characterized by dopamine deficits or cocaine abuse.

Brief Description of the Drawings

Fig. 1 is a graph depicting association of ^{125}I iodoaltropane with the dopamine transporter. The pharmacological specificity of [^{125}I] iodoaltropane labeled sites (primate striatum) and the dopamine transporter sites (rodent striatum) are closely associated.

Figs. 2A-2F are photographs depicting the distribution of ^{125}I Iodoaltropane in a primate brain, indicating a close match between that distribution and the distribution of dopamine.

Fig. 3 is a diagram of the synthesis of Iodoaltropane.

Detailed Description of the Drawings

I. Compound Characterization

Target compounds are prepared as the free bases or salts, e.g., the naphthalene-1,5-disulfonate, tartrate, or hydrochloride salts. For example, these compounds are prepared from the corresponding Z or E iodo-tri-n-butylstannylallyl precursors. See Example 1, below.

Compounds can be radiolabeled at any of a number of positions. For example, a radionuclide of the halogen series may be used in the haloallyl moiety. Alternatively, a radionuclide can be included in the -R moiety defined above. Radiolabeled iodoallyl compounds may be prepared by reacting the iodoallyl Sn precursor (see Fig. 3) with a radioiodide (provided, e.g., as NaI) under oxidative conditions (e.g., H_2O_2 or benzoyl peroxide).

- 7 -

Characterization of the compounds is carried out using standard methods of high field NMR spectra as well as IR, MS and optical rotation. Elemental analysis, TLC and/or HPLC is used as a measure of purity. A purity of >98% is preferred before biological evaluation of these compounds is undertaken.

II. Binding Assays for Candidate Compounds

To evaluate the selectivity of N-haloallyl nortropane derivatives for the dopamine transporter, compounds are screened in radioreceptor assays using both [³H]CFT and [³H]citalopram as probes for the dopamine and serotonin receptors, respectively. The relative affinities of a compound for either site establishes whether it binds selectively to the dopamine transporter or non-selectively to the serotonin transporter as well. Assays are carried out as follows.

Competition studies to determine the potency of a compound for inhibiting specifically bound [³H]CFT in monkey caudate-putamen membranes are conducted using 0.3-10nM [³H]CFT and the primate brain tissue preferably caudate-putamen. The methods are described by Madras et al. (1989) *Mol. Pharmacol.* 36:518-524. Unlabeled (-)-cocaine (30 μ M) serves as the baseline drug to detect non-specific binding. Incubation proceeds at 4°C for 60 min, and the assay is terminated by vacuum filtration over glass fiber filters (Whatman GF/B). The filters are monitored for radioactivity by liquid scintillation spectrometry at 50% counting efficiency. All assays are performed in triplicate, and each experiment is repeated at least twice using tissue from different brains.

To assay binding of the analog to the serotonin transporter, analogs are tested for their ability to compete with labeled [³H]citalopram, a high affinity and selective ligand for serotonin transporter sites (D'Amato et al., *J. Pharmacol. Exp. Ther.* 242:364-371, 1987).

- 8 -

Radioreceptor assays are conducted using tissues prepared as described above. Analogs are incubated with buffer (50 mM Tris HCl; 100mM NaCl), [³H]citalopram (1 nM), and tissue (1 mg/ml wet tissue weight), incubated 2 h at 4°C, and the incubation terminated by rapid filtration. Non-specific binding is monitored with fluoxetine (1 μM).

Using such an approach, the compounds according to the invention may be tested for their ability to inhibit the binding of [³H]CFT to primate caudate-putamen. Inhibition by (-)-cocaine and CFT may be used as reference points.

III. SPECT or PET IMAGING

Autoradiographic distribution of the compounds are conducted according to in vitro techniques (Kaufman et al. (1991) *Synapse* 9:177-187) or ex vivo techniques (Kaufman and Madras (1992) *Synapse* 12:99-111).

The cocaine analogs described herein provide useful SPECT or PET imaging probes. Brain imaging has at least two applications: to monitor the dopamine transporter (dopamine nerve terminals) in neurodegenerative disorders and cocaine abuse; and to evaluate the time course of accumulation of a candidate drug therapy in the brain targeting the dopamine transporter, as well as the duration of receptor occupancy. Such compounds may be useful for cocaine abuse or other neuro-psychiatric disorders.

Cocaine analogs of high affinity are most useful as SPECT (or PET) imaging probes because they display a low level of non-specific binding, and they accumulate in dopamine regions of brain. High affinity analogs are preferable because dopamine may compete effectively with trace doses of low affinity analogs in vivo. By these standards, Iodoaltropane is a high affinity ligand.

SPECT or PET imaging may be carried out using any appropriate apparatus. Imaging is carried out on

- 9 -

conscious human subjects using standard imaging (see, e.g., *Medicine*, Scientific American, Inc., ed. Rubenstein and Federman, 1988; Jaszczak and Coleman, *Invest. Radiol.* 20:897, 1985; Coleman et al., *Invest. Radiol.* 21:1, 1986); preferably SPECT imaging employs gamma-emitting derivatives of the analogs described herein (e.g., analogs labelled with ^{123}I).

IV. Analytical Considerations

All target compounds are prepared either as free bases or as suitable pharmacologically active salts such as hydrochloride, tartrate, naphthalene-1,5-disulfonate salts or the like. All target compounds are characterized and their purity analyzed prior to any biological evaluation. High field NMR spectra are measured as well as IR, MS and optical rotation for all test compounds. Elemental analysis, TLC and/or HPLC are used as a measure of purity. A purity of >98% is required before any biological evaluation of these compounds is undertaken.

The following specific examples illustrate but does not limit the invention.

Example 1: General Procedures for the Preparation of Iodoaltropine

Tributyltinallyl alcohol

Propargyl alcohol (5.5. mL, 9.4 mmol) was added dropwise to a mixture of tributyltin hydride (35 mL, 13 mmol) and azobis (isobutyronitrile) (AIBN) (1.6 g, 0.97 mmol) at 80°C. The reaction mixture was heated at 80°C for 2 h. The reaction mixture was chromatographed over silica gel (5% EtOAc/hexane) to afford the E-isomer 16.68g (51%) and the Z-isomer (23%) as a clear oil.

Tributyltinallyl chloride

Tributyltinallyl alcohol (12.8 g, 3.7 mmol), triphenylphosphine (9.7 g, 3.7 mmol) and CCl_4 (100 mL) were combined and heated at reflux for 16 h. Excess CCl_4 was removed and the residue was chromatographed over

- 10 -

silica gel (hexane) to afford 10.78 g (80%) of tributyltin allyl chloride as a clear oil.

Methyl-3 β -(p-fluorophenyl)-N-tributyltinallyl-1 α H-, 5 α H-nortropane-2 β -carboxylate

Methyl-3 β -(p-fluorophenyl)-1 α H, 5 α H-nortropane-2 β -carboxylate (102 g, 0.39 mmol), (see Meltzer et al., *J. Med. Chem.* **36**:855 (1993)) tributyltin allyl chloride (143 mg, 0.39 mmol), KF-celite (50%, 226 mg, 1.9 mmol) and CH₃CN (10 mL) were combined and heated at 70°C for 17 h. The reaction mixture was diluted with 40 mL of ether and filtered through a celite pad. The filtrate was concentrated to dryness. The residue was chromatographed over silica gel (2% Et₃N/hexane) to afford 163 mg (71%) of methyl-3 β -(p-fluorophenyl)-N-tributyltinallyl-1 α H-, 5 α H-nortropane-2 β -carboxylate.

Methyl-3 β -(p-fluorophenyl)-N-iodoallyl-1 α H-, 5 α H-nortropane-2 β -carboxylate

Methyl-3 β -(p-fluorophenyl)-N-tributyltinallyl-1 α H-, 5 α H-nortropane-2 β -carboxylate (161 mg, 0.27 mmol) in THF (10 mL) was degassed by bubbling N₂ through for 5 min. NIS (61.8 mg, 0.275 mmol) was added and stirred at room temperature for 15 min. THF was removed and the residue was chromatographed over silica gel (2% Et₃N/hexane) to afford 102 mg (87%) of methyl-3 β -(p-fluorophenyl)-N-iodoallyl-1 α H-, 5 α H-nortropane-2 β -carboxylate as a white solid: mp 112-114°C.

Thus, propargyl alcohol was treated with tributyltin hydride in the presence of azobis (isobutyronitrile) to obtain, after column chromatography over silica gel, the separated E and Z isomers of 3-tributyltin allyl alcohol. Each of the alcohols was then reacted with triphenylphosgene and carbon tetrachloride to obtain the requisite E and Z isomers of 3-tributyltin allyl chloride respectively.

- 11 -

Reaction of each of the 3-tributyltin allyl chlorides with methyl-3 β -(p-fluorophenyl)nortropane-2 β -carboxylate (see Meltzer et al., *J. Med. Chem.* **36**:855 (1993) for the synthesis of this compound) in the presence of KF-celite gave the desired methyl-3 β -(p-fluorophenyl)-N-(3-tributyltin allyl)-nortropane-2 β -carboxylate in 71% yield after column chromatography. Conversion of this compound to the iodide was effected with N-iodosuccinimide (NIS) in THF to obtain methyl-3 β -(p-fluorophenyl)-N-(3-iodo allyl)-nortropane-2 β -carboxylate (Iodoaltropene) in 87% yield.

Example 2: Preparation of Bromoaltropene

The 2- β -carbomethoxy-3- β -(4-fluorophenyl)-8-(3-bromopropen-2-yl) analog ("Bromoaltropene") is prepared generally as described above using N-bromosuccinimide (NBS) in place of NIS.

Example 3: Fluorination

The fluoroallyl compounds may be prepared by reacting a corresponding haloallyl Sn compound with Xe₂F₂, acetylhypofluoride, or F₂.

Example 4

Iodoaltropene was compared to CFT as well as to another candidate iodinated dopamine transporter probes known as RTI-55,² using the above-described radioreceptor binding techniques on monkey striation homogenate. The results are as follows.

1. Iodoaltropene is more selective for the dopamine transporter and possesses higher affinity than CFT (Table 1).

2. Iodoaltropene is more selective than RTI-55, which has a high affinity but low selectivity for the dopamine transporter (Table 1).

² 2-carboxymethyl-3-(4-iodophenyl)tropane.

- 12 -

3. [^{125}I]Iodoaltropine labels the dopamine transporter in primate striatum (see Fig. 1).

4. [^{125}I]Iodoaltropine distributes primarily to dopamine-rich brain regions in primate brain (Figs. 2A-2F) and its binding is characterized by low levels of nonspecific binding.


Table 1
Affinity of Iodoaltropine for the dopamine and serotonin transporter

<u>Selectivity</u>	<u>Dopamine trans.</u>	<u>Serotonin trans.</u>	
Iodoaltropine	6.62 ± 0.78	182 ± 46	28
CFT	12.9 ± 1.10	160 ± 20	15
RTI-55	1.08 ± 0.06	2.53 ± 0.02	2
Bromoaltropine	10.8	212	20

What is claimed is:

- 13 -


1. A compound of formula:



wherein the following conditions are imposed on that formula:

R is $-\text{CH}_3$, $-\text{CH}_2\text{CH}_3$, $-\text{CH}(\text{CH}_3)_2$, $-(\text{CH}_2)_n\text{CH}_3$, $-(\text{CH}_2)_n\text{C}_6\text{H}_4\text{X}$, $-\text{C}_6\text{H}_4\text{X}$, $-\text{C}_6\text{H}_5$, $-\text{OCH}_3$, $-\text{OCH}_2\text{CH}_3$, $-\text{OCH}(\text{CH}_3)_2$, $-\text{OC}_6\text{H}_5$, $-\text{OC}_6\text{H}_4\text{X}$, $-\text{O}(\text{CH}_2)_n\text{C}_6\text{H}_4\text{X}$, or $-\text{O}(\text{CH}_2)_n\text{CH}_3$; wherein X is $-\text{Br}$, $-\text{Cl}$, $-\text{I}$, $-\text{F}$, $-\text{OH}$, $-\text{OCH}_3$, $-\text{CF}_3$, $-\text{NO}_2$, $-\text{NH}_2$, $-\text{CN}$, $-\text{NHCOCH}_3$, $-\text{N}(\text{CH}_3)_2$, $-(\text{CH}_2)_n\text{CH}_3$, CHOCH_3 , or $-\text{C}(\text{CH}_3)_3$, and n is between 0 and 6 inclusive.

2. A compound of formula:



wherein the following conditions are imposed on that formula:

- 14 -

a) R is $-\text{CH}_3$, $-\text{CH}_2\text{CH}_3$, $-\text{CH}(\text{CH}_3)_2$, $-(\text{CH}_2)_n\text{CH}_3$, $-(\text{CH}_2)_nC_6H_4X$, $-\text{C}_6H_4X$, $-\text{C}_6H_5$, $-\text{OCH}_3$, $-\text{OCH}_2\text{CH}_3$, $-\text{OCH}(\text{CH}_3)_2$, $-\text{OC}_6H_5$, $-\text{OC}_6H_4X$, $-\text{O}(\text{CH}_2)_nC_6H_4X$, or $-\text{O}(\text{CH}_2)_n\text{CH}_3$; wherein X is $-\text{Br}$, $-\text{Cl}$, $-\text{I}$, $-\text{F}$, $-\text{OH}$, $-\text{OCH}_3$, $-\text{CF}_3$, $-\text{NO}_2$, $-\text{NH}_2$, $-\text{CN}$, $-\text{NHCOCH}_3$, $-\text{N}(\text{CH}_3)_2$, $-(\text{CH}_2)_n\text{CH}_3$, CHOCH_3 , or $-\text{C}(\text{CH}_3)_3$, and n is between 0 and 6 inclusive; and

b) where -Q is a halogen.

3. The compound of claim 1 or claim 2 in which said 2 substituent is in the β position.

4. The compound of claim 1 or claim 2 in which said 3 substituent is in the β position.

5. The compound of claim 1 or claim 2 in which R is $-\text{O}-\text{CH}_3$.

6. The compound of claim 1 or claim 2 in which said 8 substituent is the E isomer.

7. The compound of claim 1 or claim 2 in which said 8 substituent is the Z isomer.

8. The compound of claim 1 or claim 2 in which said compound is 2- β -carbomethoxy-3- β -(4-fluorophenyl)-8-(3-iodopropen-2-yl) nortropine.

9. A compound of claim 1 or claim 2 in which said compound contains a radioactive or fluorescent label.

10. The compound of claim 9 in which said label is a gamma or positron emitter.

- 15 -

11. The compound of claim 10 in which said label is selected from the group consisting of ^{125}I , ^{123}I , ^{18}F and ^{11}C .

12. The compound of claim 10 in which said label is ^{123}I .

13. The compound of claim 2 where Q is -Br.

14. A method of detecting parkinsonism in a human patient comprising administration to said patient of a detectably labelled compound according to claim 1 or claim 2 and detecting the binding of said compound to CNS tissue.

15. A method of selectively monitoring cocaine binding regions of the central nervous system of a human patient comprising administering to the central nervous system a detectably labelled compound according to claim 1 or claim 2 and detecting the binding of that compound to CNS tissue.

16. The method of claim 14 in which detecting is by position emission tomography (PET).

17. The method of claim 14 in which said detection is by single-photon emission computed tomography (SPECT).

FIG. 1


RELATIVE POTENCY
DOPAMINE UPTAKE

FIG. 2A



B




FIG. 2D

D



FIG. 2C

C



FIG. 2F

F





FIG. 2E

E



3/3

FIG. 3



INTERNATIONAL SEARCH REPORT

International application No.
PCT/US94/12225

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : CO7D 451/02; A61K 51/02, 101/02, 31/41

US CL : 546/132; 514/304; 424/1.85, 1.89

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 546/132; 514/304; 424/1.85, 1.89

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
STN - Structure Search

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A, P	US, A, 5,268,480 (KOZIKOWSKI) 07 DECEMBER 1993, See abstract	1-8
Y	JOURNAL OF NUCLEAR MEDICINE, Vol. 33(5), May 1992 GOODMAN ET AL., "Radioiodinated 2beta - Carbomethoxy - 3beta (4-Chlorophenyl)-8-(3E-and 3-Z-Iodopropen-2-yl) NORTROPANES....," Abstract # 275, page 890, see figure	1-17
Y	PHARMACOLOGY, BIOCHEMISTRY & BEHAVIOR, Vol. 35, 1990, MADRAS ET AL. "N-Modified Fluorophenyltropene Analogs ...", pp. 949-953, see page 951.	1-17
Y	JOURNAL OF MEDICINAL CHEMISTRY, VOL. 34(9), 1991, CARROLL ET AL., "Synthesis, Ligand Binding, QSAR, and CoMFA Study....," pp. 2719-2725, see Table I & II	1-17

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:

A document defining the general state of the art which is not considered to be part of particular relevance

E earlier document published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T

later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X

document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y

document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

&

document member of the same patent family

Date of the actual completion of the international search

23 FEBRUARY 1995

Date of mailing of the international search report

23 FEB 1995

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

GARY GEIST

Telephone No. (703) 308-0450

Form PCT/ISA/210 (second sheet)(July 1992)*

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US94/12225

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	JOURNAL OF MEDICINAL CHEMISTRY, VOL. 34(4) 1991, MILIUS ET AL., "Synthesis and Receptor Binding of N-Substituted Tropane Derivatives...", pp. 1728-1731, see Table I.	1-17